

VIRIYA

energy / virility

PROPRIETARY NAME AND DOSAGE FORM:

VIRIYA capsules

COMPOSITION:

Each capsule provides:

Tongkat ali (<i>Eurycoma longifolia</i>)	138,60 mg
Hong hua (<i>Flos cathami</i>)	22,05 mg
Chiang huang (<i>Rhizoma curcuma longa</i>)	44,10 mg
Bai gou ye (<i>Ginkgo biloba</i>)	44,10 mg
Yin yang huo (<i>Herba epimedii</i>)	22,05 mg
Rou cong rong (<i>Herba cistanches</i>)	22,05 mg
Huang qi (<i>Astragalus membranaceus</i>)	22,05 mg
Gin seng (<i>Ren shen</i>)	25,00 mg
Gou qi zi (<i>Fructus lycii</i>)	75,00 mg
Gan cao (<i>Liquorice</i>)	25,00 mg
Sugar free.	

PHARMACOLOGICAL CLASSIFICATION:

A: 32,16 Others

INDICATIONS:

Provides energy, improves virility.

CONTRA-INDICATIONS:

Do not use excessive panax ginseng with pre-diagnosed high blood pressure, or if calcium deficient, except under practitioner supervision.

WARNINGS:

Seek professional advice before using long term.

INTERACTIONS:

Do not use ginkgo biloba with warfarin except under practitioner supervision.

PREGNANCY AND LACTATION:

Do not use panax ginseng if pregnant.

DOSAGE AND DIRECTIONS FOR USE:

Take one capsule with a little water an hour before an intimate encounter. Some users may find two capsules more effective. The benefits of the product should be felt for between 12 and 30 hours. Unless otherwise prescribed, do not exceed the stated dose.



SIDE EFFECTS AND SPECIAL PRECAUTIONS:

Ginkgo biloba may cause headaches, dizziness, palpitations, gastro-intestinal disturbances, bleeding disorders and skin hypersensitivity reactions. (Ref 1)

Astragalus may cause hypersensitivity reactions. (Ref 2)

Panax ginseng may cause diarrhoea, skin eruptions, sleeplessness, nervousness, hypertension, euphoria, oedema, oestrogenic effects, and rarely Stevens-Johnson syndrome. (Ref 3)

Liquorice may cause sodium and water retention, and hypokalaemia. (Ref 4)

Caution: Rhizoma curcuma longa in large doses may cause increased sensitivity to sunlight especially if the person is fair skinned. Since chaparral may cause liver damage, use only under practitioner supervision.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

None known.

IDENTIFICATION:

Clear capsules containing beige powder.

PRESENTATION:

Blister strips of 2's, 5's and 10's in clear pvc on silver foil, packed in cardboard cartons.

STORAGE INSTRUCTIONS:

Store below 30°C in a cool dry place, out of direct sunlight.

Keep out of reach of children.

NAME AND BUSINESS ADDRESS OF THE APPLICANT:

INDIGO TRADING (PTY) LTD.

P.O. Box 785100, Sandton, 2146.

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

June 2007

REFERENCES:

1. Martindale, 33rd edition, page 1615.2

2. Martindale, 33rd edition, page 1504.2

3. Martindale, 33rd edition, page 1615.3

4. Martindale, 33rd edition, page 1231.2

